

A negative answer to a one-dimensional symmetry problem in the Heisenberg group

I. Birindelli*, E. Lanconelli†

1 Introduction and Main Results

Symmetry properties of solutions to semi-linear elliptic equations have been widely studied in the last decades. In this contest, a long standing conjecture by De Giorgi states that any global solution to the Ginzburg-Landau equation

$$\Delta u + u(1 - u^2) = 0 \text{ in } \mathbb{R}^N \text{ } (N \leq 8)$$

satisfying $-1 \leq u \leq 1$ and $\frac{\partial u}{\partial x_N} > 0$ is constant along hyperplanes. Recently this conjecture was proved to be true by Ghoussoub and Guy for $N = 2$ ([15]) and by Ambrosio and Cabré for $N = 3$ ([3]). It is still an open question for $N > 3$ though Albriti, Ambrosio and Cabré generalized the result for any C^2 non-linearity (when $N \leq 3$) [1].

Under the further hypothesis that the solution u satisfies

$$\lim_{x_3 \rightarrow \pm\infty} u(x', x_3) = \pm 1 \quad \forall x' \in \mathbb{R}^2$$

the proof that u is constant along hyperplanes given in [3] is somehow simpler. On the other hand, under the hypothesis that this limit is uniform in x' , the conjecture was known as Gibbons conjecture and it has been proved for all dimensions independently by Barlow, Bass, Guy in [4], Berestycki, Hamel, Monneau in [5] and Farina in [11].

*Dipartimento di Matematica, Università di Roma "La Sapienza", Piazzale A.Moro 2 - I 00185 Roma, Italy; e-mail: isabeau@mat.uniroma1.it

†Dipartimento di Matematica, Università di Bologna, Piazza di Porta San Donato 5, I 40126 Bologna; e-mail: lanconelli@mat.unibo.it

In recent years symmetry and monotonicity properties of solutions to semilinear equations have been investigated in the more general context of the Carnot groups, see [7, 8, 9], [2], [6] and [13]. The interest in semi-linear equations in these groups has increased as they appear in many theoretical and application fields, such as complex geometry and mathematical models for crystal structures [10].

In [8] Prajapat and the first author studied Gibbons conjecture for the equation

$$\Delta_{\mathbb{H}^n} u + f(u) = 0 \text{ in } \mathbb{H}^n, \quad (1.1)$$

where $\Delta_{\mathbb{H}^n}$ denotes the Kohn-Laplacian on the Heisenberg group $\mathbb{H}^n$ and $f(u)$ is a non linear term with some general hypothesis (in particular they include the case $f(u) = u(1 - u^2)$). They prove that the conjecture holds true for all directions orthogonal to the center of $\mathbb{H}^n$.¹ The question of whether the result holds true in the remaining direction was raised in [8].

The aim of this paper is to prove that, with respect to the center direction of $\mathbb{H}^n$, the stronger De Giorgi conjecture is not true for the equation (1.1). This negative answer will easily follow from next Theorem 1.1, the main result of this note.

In order to clearly state our theorem, we need to recall some known facts about the Heisenberg space $\mathbb{H}^n$ and its intrinsic Laplacian $\Delta_{\mathbb{H}^n}$.

First of all let us say that $\mathbb{H}^n$ is the Lie group whose underlying manifold is $\mathbb{C}^n \times \mathbb{R}$, $n \in \mathbb{N}$, endowed with the group action $\circ$ given by

$$\xi_o \circ \xi = (z + z_o, t + t_o + 2\text{Im}(\bar{z} \cdot z_o)). \quad (1.2)$$

Here and in the rest of the paper we identify $\mathbb{C}^n$ with $\mathbb{R}^{2n}$ and, setting $z = x + iy$, for the point of $\mathbb{H}^n$ we use the equivalent notations $\xi = (z, t) = (x, y, t) \in \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$ with $z := (z_1, \dots, z_n) = (x_1, y_1, \dots, x_n, y_n)$. Furthermore, “ $\cdot$ ” denotes the usual inner product in $\mathbb{C}^n$.

The Lie Algebra of left invariant vector fields is generated by

$$\begin{aligned} X_i &= \frac{\partial}{\partial x_i} + 2y_i \frac{\partial}{\partial t}, \quad \text{for } i = 1, \dots, n, \\ Y_i &= \frac{\partial}{\partial y_i} - 2x_i \frac{\partial}{\partial t}, \quad \text{for } i = 1, \dots, n. \end{aligned}$$

The intrinsic Laplacian of $\mathbb{H}^n$, also called the Kohn Laplacian, is defined as

$$\Delta_{\mathbb{H}^n} = \sum_{i=1}^n (X_i^2 + Y_i^2).$$

¹Very recently, in [6], the results of [8] have been extended to every sub-Laplacian on a Carnot group.

It is a second order degenerate elliptic operator of Hormander type and hence it is hypoelliptic (see e.g. [12] or [16] for more details about $\Delta_{\mathbb{H}^n}$).

With respect to the group dilation $\delta_\lambda \xi = (\lambda z, \lambda^2 t)$, $\Delta_{\mathbb{H}^n}$ is homogeneous of degree two in the following sense

$$\Delta_{\mathbb{H}^n} \circ \delta_\lambda = \lambda^2 \delta_\lambda \circ \Delta_{\mathbb{H}^n}.$$

The Koranyi ball of center ξ_o and radius R is defined by

$$B_H(\xi_o, R) := \{\xi \text{ such that } |\xi^{-1} \circ \xi_o|_{\mathbb{H}^n} \leq R\}$$

where

$$|\xi|_{\mathbb{H}^n} = (|z|^4 + t^2)^{\frac{1}{4}}$$

is a norm with respect to the group dilation and it satisfies

$$|B_H(\xi_o, R)| = |B_H(0, R)| = CR^Q$$

where $Q = 2n + 2$ is the homogeneous dimension of $\mathbb{H}^n$.

A fundamental solution of $-\Delta_{\mathbb{H}^n}$ with pole at the origin is given by:

$$\Gamma(\xi) = \frac{C_Q}{(|\xi|_{\mathbb{H}^n})^{Q-2}}$$

where C_Q is a positive constant.

For our purposes it is convenient to remind that the class of cylindrically symmetric functions is invariant with respect to the action of $\Delta_{\mathbb{H}^n}$. We shall say that a function $(z, t) \rightarrow u(z, t)$ is cylindrically symmetric if there exists a two variables function U such that $u(z, t) = U(r, t)$, $r = |z|$.

In that case we formally have that

$$\Delta_{\mathbb{H}^n} u(z, t) = \partial_{rr} U + \frac{2n-1}{r} \partial_r U + 4r^2 \partial_{tt} U.$$

The main result of this paper is the following:

Theorem 1.1 *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a locally Lipschitz continuous function satisfying the hypotheses listed below:*

- (H1) *f is odd,*
- (H2) *$f > 0$ in $]0, 1[$, $f(0) = f(1) = 0$,*
- (H3) *$\lim_{s \rightarrow 0} \frac{f(s)}{s} = l > 0$.*

Then there exists a solution u to the equation:

$$\Delta_{\mathbb{H}^n} u + f(u) = 0 \quad \text{in } \mathbb{R}^{2n+1} \quad (1.3)$$

satisfying $|u| < 1$, $\frac{\partial u}{\partial t} > 0$ and

$$\lim_{t \rightarrow \pm\infty} u(z, t) = \pm 1.$$

Moreover u is cylindrically symmetric and of class C^∞ when f is C^∞ .

For solution u of (1.3) we mean a continuous function u such that:

1. For a suitable $\alpha > 0$, $u \in \Lambda_{\text{loc}}^{2+\alpha}(\mathbb{H}^n)$ i.e. $X_j^2 u$ and $Y_j^2 u$, $j = 1, \dots, n$, exists in the weak sense of distributions and belong to $\Lambda_{\text{loc}}^\alpha(\mathbb{H}^n)$
2. u satisfies (1.3) pointwise everywhere.

As in [12] we have denoted by $\Lambda_{\text{loc}}^\alpha(\mathbb{H}^n)$ the class of functions which are locally α -Holder continuous with respect to the intrinsic distance d in $\mathbb{H}^n$ defined by

$$d(\xi, \xi') = |(\xi')^{-1} \circ \xi|_{\mathbb{H}^n}.$$

Using the commutators of the Lie Algebra, it is easy to see that $\Lambda_{\text{loc}}^{2+\alpha}(\mathbb{H}^n)$ is continuously embedded in the usual $C_{\text{loc}}^{1+\frac{\alpha}{2}}(\mathbb{R}^{2n+1})$.

From Theorem 1.1 we immediately get the following corollary.

Corollary 1.1 *De Giorgi's conjecture in the t -direction is not true in $\mathbb{H}^n$.*

Proof. The functions $f(s) = s(1-s^2)$ satisfies all hypotheses of Theorem 1.1, hence there exists a C^∞ function u such that

$$\begin{cases} \Delta_{\mathbb{H}^n} u + u(1-u^2) = 0 & \text{in } \mathbb{R}^{2n+1}, \\ -1 < u < 1, \frac{\partial u}{\partial t} > 0, \\ \lim_{t \rightarrow \pm\infty} u(z, t) = \pm 1. \end{cases}$$

Then, if De Giorgi conjecture were true in the t direction there would exist $\alpha \in \mathbb{R}^{2n}$ and $\nu > 0$ such that $u(z, t) = U(\alpha \cdot z + t\nu)$ for some function $U : \mathbb{R} \rightarrow \mathbb{R}$. Furthermore U would satisfy

$$(|\alpha|^2 - 4\nu(J\alpha \cdot z) + 4r^2\nu^2) U'' = U(U^2 - 1)$$

where J is the classical symplectic $2n \times 2n$ matrix. This is a contradiction since the right hand side is constant along the hyperplanes $\alpha \cdot z + t\nu = c$ for any $c \in \mathbb{R}$ while the left hand side is not.

Remark 1.1 It would be interesting to know whether the function constructed in Theorem 1.1 has uniform limit with respect to z .

Remark 1.2 It is natural to consider the extension of Theorem 1.1 to the contest of Carnot groups. This will be the object of a subsequent study.

2 Proof of Theorem 1.1.

For any $R > 0$ we shall denote by D_R and D_R^+ respectively the cylinders

$$D_R = \{(z, t) \in \mathbb{R}^{2n+1}; |z| < R, |t| < R^2\}$$

and

$$D_R^+ = \{(z, t) \in \mathbb{R}^{2n+1}; |z| < R, 0 < t < R^2\}.$$

Let $\psi(t) = \frac{t}{R^2}$.

We shall split the proof in several steps.

First step: *The semilinear Dirichlet problem*

$$\begin{cases} \Delta_{\mathbb{H}^n} u = -f(u) & \text{in } D_R^+, \\ u(r, t) = \psi(t), & \text{on } \partial D_R^+. \end{cases} \quad (2.4)$$

has a solution $u \in \Lambda_{\text{loc}}^{2+\alpha}(D_R^+) \cap \Lambda^\alpha(\overline{D_R^+})$ for a suitable $\alpha \in (0, 1)$. Furthermore u is cylindrically symmetric, $0 \leq u \leq 1$ and for any R sufficiently large,

$$u \geq v_o$$

for some function $v_o \geq 0$, $v_o \not\equiv 0$, v_o independent of R .

Let $M \in \mathbb{R}^+$ be larger than the Lipschitz constant of f in $[0, 1]$ and let us define

$$g : \mathbb{R} \rightarrow \mathbb{R}, \quad g(s) = f(s) + Ms.$$

Let $\mathcal{T}$ be the map formally defined by $\mathcal{T}(v) = u$ where u is the only solution to the Dirichlet problem

$$\begin{cases} \Delta_{\mathbb{H}^n} u - Mu = -g(v) & \text{in } D_R^+, \\ u = \psi, & \text{on } \partial D_R^+. \end{cases} \quad (2.5)$$

The operator $\mathcal{T}$ has the following properties:

(P1) There exists $\alpha \in (0, 1)$ such that $\mathcal{T}$ is well defined in $\Lambda^\alpha(\overline{D_R^+})$. Furthermore

$$|u(\xi) - u(\xi')| \leq Cd(\xi', \xi)^\alpha (1 + \sup |g(v)|) \quad (2.6)$$

for any $\xi, \xi' \in D_R^+$. We also have that $\mathcal{T}(v) \in \Lambda_{\text{loc}}^{2+\alpha}(D_R^+)$ for every $v \in \Lambda^\alpha(\overline{D_R^+})$.

This statement can be proved by using standard arguments and the results in [12], [16] (see also [14, Theorem 4.1]).

(P2) $\mathcal{T}(v)$ is cylindrically symmetric if v is cylindrically symmetric.

Indeed suppose that $u = \mathcal{T}(v)$. Let $\mathcal{S}$ be a rotation in $\mathbb{R}^{2n}$ and define $u_{\mathcal{S}}(z, t) := u(\mathcal{S}z, t)$. Since $\Delta_{\mathbb{H}^n}$ is invariant with respect to $\mathcal{S}$, we have $\Delta_{\mathbb{H}^n} u_{\mathcal{S}}(z, t) = \Delta_{\mathbb{H}^n} u(\mathcal{S}z, t)$, so that $u_{\mathcal{S}}$ is a solution of

$$\begin{cases} \Delta_{\mathbb{H}^n} u_{\mathcal{S}} - M u_{\mathcal{S}} = -g(v(\mathcal{S}z, t)) = -g(v) & \text{in } D_R^+, \\ u_{\mathcal{S}} = \psi, & \text{on } \partial D_R^+. \end{cases}$$

Here we have used the invariance with respect to $\mathcal{S}$ of v , ψ and D_R^+ .

By the maximum principle we know that the solution of (2.5) is unique, hence $u = u_{\mathcal{S}}$ for any $\mathcal{S}$, i.e. u is cylindrically symmetric.

(P3) $\mathcal{T}$ is monotone. More precisely if $v_1, v_2 \in \Lambda^\alpha(\overline{D_R^+})$ and $0 \leq v_1 \leq v_2 \leq 1$, then $\mathcal{T}v_1 \leq \mathcal{T}v_2$.

Let us observe that with our choice of M if $0 \leq v_1 \leq v_2$ then $g(v_1) \leq g(v_2)$. Hence (P3) follows from the maximum principle for $-\Delta_{\mathbb{H}^n} + M$ in D_R^+ .

(P4) If $v \in \Lambda^\alpha(\overline{D_R^+})$ and $0 \leq v \leq 1$ then $0 \leq \mathcal{T}(v) \leq 1$.

Indeed, since $g(0) = 0$, $g(1) = M$ and $0 \leq \psi \leq 1$ on ∂D_R^+ , again by the maximum principle we obtain that $\mathcal{T}(1) \leq 1$ and $\mathcal{T}(0) \geq 0$. Now we only need to apply property (P3) for $v \in \Lambda^\alpha(\overline{D_R^+})$ such that $0 \leq v \leq 1$.

We shall now construct a function $v_o \geq 0$ that plays the role of a lower barrier.

Let λ_o denote the principal eigenvalue of $-\Delta_{\mathbb{H}^n}$ in D_R^+ and let $\phi_o > 0$ be the corresponding eigenfunction normalized by $\sup \phi_o = 1$.

We choose and fix R_o sufficiently large that

$$\lambda_o \leq \frac{l}{2}$$

where l is the limit in condition (H3). Then there exists $\varepsilon \in (0, 1)$ independent of R such that

$$\lambda_o \varepsilon \phi_o \leq f(\varepsilon \phi_o).$$

By uniqueness of the normalized eigenfunction ϕ_o , arguing as in the proof of (P2) we can prove that ϕ_o is cylindrically symmetric.

From now on we assume that $R > R_o$. Let us define

$$v_o = \begin{cases} \varepsilon \phi_o & \text{in } D_{R_o}^+ \\ 0 & \text{in } D_R^+ \setminus D_{R_o}^+. \end{cases}$$

Standard arguments show that v_o is locally Holder continuous in $\mathbb{R}^{2n+1}$, (see e.g. [14, Theorem 4.1], we stress that condition (4.4) in that theorem is satisfied since D_R^+ is convex).

As a consequence $\mathcal{T}(v_o)$ is well defined and since $0 \leq v_o \leq 1$ using (P4) we get that $0 \leq \mathcal{T}(v_o) \leq 1$. Let us now prove that $v_o \leq u_o := \mathcal{T}(v_o)$. Clearly the inequality holds in $D_R^+ \setminus D_{R_o}^+$, using (P4), hence we just have to prove it in $D_{R_o}^+$. We have

$$\begin{aligned} \Delta_{\mathbb{H}^n} u_o - M u_o &= -g(v_o) = -g(\varepsilon \phi_o) \leq -(M + \lambda_o)(\varepsilon \phi_o) \\ &= -M \varepsilon \phi_o + \Delta_{\mathbb{H}^n} \varepsilon \phi_o = -M v_o + \Delta_{\mathbb{H}^n} v_o, \end{aligned}$$

so that

$$\begin{cases} \Delta_{\mathbb{H}^n}(u_o - v_o) - M(u_o - v_o) \leq 0 & \text{in } D_{R_o}^+ \\ u_o \geq v_o & \text{on } \partial D_{R_o}^+. \end{cases}$$

The maximum principle implies that $u_o \geq v_o$ in $D_{R_o}^+$.

Now we construct the sequence of functions

$$v_k = \mathcal{T}^k(v_o), k \in \mathbb{N}.$$

Clearly using the properties above, all v_k are cylindrically symmetric and

$$1 \geq \mathcal{T}^k(v_o) \geq \mathcal{T}(v_o) \geq v_o \geq 0 \text{ for every } k \in \mathbb{N}.$$

Let us denote by u the pointwise limit of (v_k) . Then u is cylindrically symmetric, $v_o \leq u \leq 1$, $u \in \Lambda^\alpha(\overline{D_R^+})$ since, by (2.6)

$$|v_k(\xi) - v_k(\xi')| \leq C d(\xi', \xi)^\alpha$$

where $C > 0$ is independent of R . This estimates implies that the v_k uniformly converges to u in $\overline{D_R^+}$, so that $u = \psi$ on ∂D_R^+ .

Furthermore in the weak sense of distributions, u satisfies

$$\Delta_{\mathbb{H}^n} u + f(u) = 0 \quad \text{in } D_R^+. \quad (2.7)$$

From (2.7), the Holder regularity of u and standard bootstrap argument we obtain that $u \in \Lambda_{\text{loc}}^{2+\alpha}(D_R^+)$ and it satisfies the equation pointwise. Hence u is the required function.

Remark 2.1 Since u is cylindrically symmetric we have that $u(z, t) = U(|z|, t)$ and U satisfies the semilinear elliptic equation

$$\partial_{rr} U + \frac{2n-1}{r} \partial_r U + 4r^2 \partial_{tt} U + f(U) = 0$$

in the open subset of $\mathbb{R}^2$

$$\Omega_R := \{(r, t) \in \mathbb{R}^2 / 0 < r < R, 0 < t < R^2\}.$$

Moreover U is locally $\frac{\alpha}{2}$ -Holder continuous, in the usual sense, up to $\partial\Omega_R \setminus \{(0, t) / 0 < t < R^2\}$. Then, being $U(r, 0) = 0$ when $0 < r < R$, by classical regularity results for elliptic equations, U is of class $C_{\text{loc}}^{2+\frac{\alpha}{2}}$ up to $\Omega_R \cup \{(r, 0) / 0 < r < R\}$.

Second step: *The function constructed in the first step satisfies $\frac{\partial u}{\partial t} > 0$.*

In [9] the following definition and theorem are given:

Definition 2.1 Fix $\eta \in \mathbb{H}^n$. A domain $\Omega \subset H$ is said to be η -convex (or convex in the direction η) if for any $\xi_1 \in \Omega$ and any $\xi_2 \in \Omega$ such that $\xi_2 = \alpha\eta \circ \xi_1$ for some $\alpha > 0$, we have $s\eta \circ \xi_1 \in \Omega$ for every $s \in (0, \alpha)$.

Theorem 2.1 Let Ω be an arbitrary bounded domain of $\mathbb{H}^n$ which is η -convex for some $\eta \in H$. Let $u \in S_2^Q(\Omega) \cap C(\bar{\Omega})$ be a solution of

$$\left. \begin{aligned} \Delta_{\mathbb{H}^n} u + f(u) &= 0 \quad \text{in } \Omega \\ u &= \phi \quad \text{on } \partial\Omega \end{aligned} \right\} \quad (2.8)$$

where f is a Lipschitz continuous function. Assume that for any $\xi_1, \xi_2 \in \partial\Omega$, such that $\xi_2 = \alpha\eta \circ \xi_1$ for some $\alpha > 0$, we have for each $s \in (0, \alpha)$

$$\phi(\xi_1) < u(s\eta \circ \xi_1) < \phi(\xi_2) \quad s\eta \circ \xi_1 \in \Omega \quad (2.9)$$

and

$$\phi(\xi_1) < \phi(s\eta\xi_1) < \phi(\xi_2) \quad \text{if } s\eta \circ \xi_1 \in \partial\Omega \quad (2.10)$$

Then u satisfies

$$u(s_1\eta \circ \xi) < u(s\eta \circ \xi) \quad (2.11)$$

for any $0 < s_1 < s < \alpha$ and for every $\xi \in \Omega$.

Moreover, u is the unique solution of (2.8) in $S_2^Q(\Omega) \cap C(\bar{\Omega})$ satisfying (2.9).

Let us choose $\eta = (0, 1)$, clearly D_R^+ is η -convex since:

$$s\eta \circ \xi = (z, t + s).$$

Furthermore $0 = \psi(0) \leq u(z, t) \leq \psi(1) = 1$ and by construction ψ satisfies (2.10). Hence we are in the hypothesis of Theorem 2.1 and u satisfies

$$u(z, t_1) \leq u(z, t_2) \quad \text{for any } 0 \leq t_1 \leq t_2 \leq 1$$

in D_R^+ .

In particular we get $\frac{\partial u}{\partial t} \geq 0$.

Now since $\frac{\partial}{\partial t}$ commutes with $\Delta_{\mathbb{H}^n}$ and f is Lipschitz continuous then the inequality is strict, just by using the strong Maximum principle.

Third step. We extend to D_R the function u of the previous step by setting

$$v(z, t) = \begin{cases} u(z, t) & \text{for } t \geq 0 \\ -u(z, -t) & \text{for } t \leq 0. \end{cases}$$

Obviously v is cylindrically symmetric, $-1 \leq v \leq 1$, $v \geq v_o$ in D_R^+ , $v \in C^{\frac{\alpha}{2}}(D_R)$ and $v = \phi$ on ∂D_R . We want to prove that v satisfies

$$\Delta_{\mathbb{H}^n} v + f(v) = 0 \quad \text{in } D_R. \quad (2.12)$$

Since f is odd, using the fact that v is odd and cylindrically symmetric it is easy to see that v satisfies (2.12) in $D_R \setminus \{t = 0\}$.

By Remark 2.1 at the end of the first step, we now obtain that $v \in C^{2+\frac{\alpha}{2}}(D_R \setminus \{(0, 0)\})$ and it solves (2.12) in the same open set. Hence we just have to remove the singularity at the origin. Let us define

$$w(\xi) = - \int_{D_R} \Gamma((\xi')^{-1} \circ \xi) f(v(\xi')) d\xi',$$

where $\Gamma(z, t)$ is the fundamental solution recalled in the Introduction. Since $f(v) \in C^{\frac{\alpha}{2}}(D_R)$ and $C^{\frac{\alpha}{2}}_{loc}(D_R) \subset \Lambda^{\frac{\alpha}{2}}_{loc}(D_R)$, then $w \in \Lambda^{2+\frac{\alpha}{2}}_{loc}(D_R)$ and satisfies

$$\Delta_{\mathbb{H}^n} w = f(v) \text{ in } D_R.$$

Hence

$$\Delta_{\mathbb{H}^n}(v + w) = 0 \text{ in } D_R \setminus \{(0, 0)\}.$$

On the other hand $v + w \in L^\infty(D_R)$. Then there exists a C^∞ -function h , $\Delta_{\mathbb{H}^n}$ harmonic in D_R such that

$$h = v + w \text{ in } D_R \setminus \{(0, 0)\}.$$

It follows that v solves (2.12) everywhere in D_R .

This ends the third step. We shall denote $u_R(z, t) = v(z, t)$ the function constructed above.

Fourth step. *We let R tend to infinity and obtain a global solution.*

Since the functions u_R are equi-bounded and solutions of (2.4) in D_R , then $\Delta_{\mathbb{H}^n} u_R$ are also equi-bounded and by standard arguments, eventually passing to a subsequence, the u_R 's locally uniformly converge to u , weak solution of

$$\Delta_{\mathbb{H}^n} u + f(u) = 0 \text{ in } \mathbb{R}^{2n+1}. \quad (2.13)$$

Furthermore

- 1) u is cylindrically symmetric,
- 2) $-1 \leq u \leq 1$,
- 3) $u(z, t) = -u(z, -t)$,
- 4) for $t \geq 0$, $u(z, t) \geq v_o(z, t)$,
- 5) $t \mapsto u(z, t)$ is monotone increasing.

Since f is locally Lipschitz continuous and $|u| \leq 1$, it follows from (2.13) that $u \in \Lambda^{2+\alpha}_{loc}(\mathbb{H}^n)$ for every $\alpha < 1$. Obviously, the more regular f is, the more regular u is; in particular u is of class C^∞ when f is C^∞ .

Moreover, property 5) implies $\frac{\partial u}{\partial t} \geq 0$ so that, since $\frac{\partial}{\partial t}$ commutes with $\Delta_{\mathbb{H}^n}$, by the strong maximum principle either $\frac{\partial u}{\partial t} > 0$ or $\frac{\partial u}{\partial t} \equiv 0$. But by 3) and 4) this second possibility is absurd hence $\frac{\partial u}{\partial t} > 0$.

Last step. *We want to prove that*

$$\lim_{t \rightarrow \pm\infty} u(z, t) = \pm 1.$$

We shall consider only the limit in $+\infty$ since the other case follows similarly. Let us denote $u_o(z) := \lim_{t \rightarrow +\infty} u(z, t)$. Since u is bounded and monotone in t the limit is well defined and $0 < u_o(z) \leq 1$. We want to prove that $u_o(z) \equiv 1$.

By standard arguments (multiplying equation (2.13) by a sequence of functions $\psi_k(z, t) = \phi(z)\phi_k(t)$ where ϕ has compact support and $\text{supp}\phi_k =]k, k+1[$ and $\int \phi_k dt = 1$ and letting k go to infinity) it is easy to see that u_o is a weak solution of

$$\Delta u_o + f(u_o) = 0 \quad \text{in } \mathbb{R}^{2n}.$$

Clearly a bootstrap argument shows that u_o is a classical solution. Moreover $u_o(z) = U_o(r)$ with $r = |z|$ for some function U_o solution of

$$U_o''(r) + \frac{2n-1}{r}U_o'(r) + f(U_o(r)) = 0, \quad (2.14)$$

$$U_o'(0) = 0 \quad (2.15)$$

The Cauchy problem for (2.14) with initial conditions $U_o(0) = 1$ and $U_o'(0) = 0$ has a unique solution (see e.g. [17]). Thus, since $f(1) = 0$, if $U_o(0) = 1$ then $U_o \equiv 1$ and we are done. Suppose, by contradiction, that $U_o(0) < 1$.

It is easy to see that $U_o' < 0$. Indeed integrating (2.14) one obtains:

$$r^{2n-1}U_o'(r) = - \int_0^r \rho^{2n-1}f(U_o(\rho))d\rho < 0, \quad (2.16)$$

hence U_o is strictly decreasing and has a finite non-negative limit as $r \rightarrow \infty$. More precisely $\lim_{r \rightarrow +\infty} U_o(r) = 0$. Indeed otherwise $U_o(r) \rightarrow k > 0$ and $f(U_o(r)) \rightarrow f(k) > 0$ (by (H3)). This, together with (2.16) implies that $|U_o'(r)| \rightarrow \infty$, which is absurd since U_o is bounded. Using hypothesis (H4) on f we obtain that for r large U_o satisfies

$$U_o''(r) + \frac{2n-1}{r}U_o'(r) + K(r)U_o(r) = 0$$

with $K(r) = \frac{f(U_o(r))}{U_o(r)} \rightarrow l > 0$.

Using the substitution $V_o(r) = r^{\frac{2n-1}{2}}U_o(r)$ we obtain that V_o satisfies

$$V_o''(r) + H(r)V_o(r) = 0$$

with $H(r) = \frac{2n-1}{2}(1 - \frac{N-1}{2})\frac{1}{r^2} + K(r)$. Comparing with

$$U''(r) + \frac{l}{2}U(r) = 0$$

we obtain that V_o i.e. U_o has infinite zeros in a neighborhood of infinity, which is absurd. This conclude the last step and the proof.

References

- [1] G. Alberti, L. Ambrosio, X. Cabré, on a long-standing conjecture of E. De Giorgi: old and recent results *Acta Applic. Math.* to appear.
- [2] L. Almeida, Y. Ge, Symmetry results for positive solutions of some elliptic equations on manifolds, *Annals of Global Analysis and geometry* to appear.
- [3] L. Ambrosio, X. Cabré, Entire solutions of semilinear elliptic equations in $\mathbb{R}^3$ and a conjecture of De Giorgi, *J. Amer. Math. Soc.* to appear .
- [4] M.T. Barlow, R. F. Bass, C. Gui, The Liouville property and a conjecture of De Giorgi, *Comm. Pure Appl. Math.***53** (2000), no. 8, 1007–1038.
- [5] H. Berestycki, F. Hamel, R. Monneau, One-dimensional symmetry of bounded entire solutions of some elliptic equations, *Duke Math. J.* **103** (2000), no. 3, 375–396.
- [6] I. Birindelli, E. Lanconelli, A note on one dimensional symmetry in Carnot Groups, *Rendiconti dell'Accademia dei Lincei* to appear.
- [7] I. Birindelli, J. Prajapat, Nonlinear Liouville theorems in the Heisenberg group via the moving plane method *Comm. Partial Differential Equations* **24** (1999)1875–1890.
- [8] I. Birindelli, J. Prajapat, One dimensional symmetry in the Heisenberg group, to appear in *Ann. Scuola Normale Superiore di Pisa*.
- [9] I. Birindelli, J. Prajapat, Monotonicity results for Nilpotent Stratified Groups, *Pacific J.* to appear .

- [10] D. Christodoulou: On the Geometry and dynamics of crystalline continua, *Ann.Inst. H.Poincaré', Phys. Theor.* **69** (1998) 335-358
- [11] A. Farina, Symmetry for solutions of semilinear elliptic equations in $\mathbb{R}^N$ and related conjectures, *Ricerche di Matematica* **XLVIII**, (1999), 129-154.
- [12] G. B. Folland, E.M. Stein, *Estimates for the ∂_b complex and analysis on the Heisenberg group*, Indiana Univ. Math. J. 41 (1992), 71-97.
- [13] N. Garofalo, D. Vassilev, Symmetry properties of positive entire solutions of Yamabe-type equations on groups of Heisenberg type, *Duke Math. J.*, **106** (2001), 411-448.
- [14] N. Garofalo, E. Lanconelli, Existence and Nonexistence Results for Semilinear Equations on the Heisenberg Group, *Indiana Univ. Math. J.*, **41** (1992), 71-98.
- [15] N. Ghoussoub, C. Gui, On a conjecture of De Giorgi and some related problems, *Math. Ann.*, **311** (1998), 481-491.
- [16] D.S. Jerison, Boundary regularity in the Dirichlet problem for b on CR manifolds *Comm. Pure Appl. Math* **36** (1983), 143-181.
- [17] L.A. Peletier, J. Serrin, Uniqueness of positive solution to semilinear equations in $\mathbb{R}^n$, *J. Differential Equations* , **61** (1986), 380-397.